

**European Community Directive
on the Conservation of Natural Habitats
and of Wild Fauna and Flora
(92/43/EEC)**

**Fourth Report by the United Kingdom
under Article 17**

on the implementation of the Directive
from January 2013 to December 2018

Conservation status assessment for the species:

S2016 - Kuhl's Pipistrelle (*Pipistrellus kuhlii*)

UNITED KINGDOM

IMPORTANT NOTE - PLEASE READ

- The information in this document represents the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- It is based on supporting information provided by the geographically-relevant Statutory Nature Conservation Bodies, which is documented separately.
- The 2019 Article 17 UK Approach document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Maps showing the distribution and range of the species are included (where available).
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the UK assessments. Further underpinning explanatory notes are available in the related country-level reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species).
- The UK-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, <https://jncc.gov.uk/article17>, for further information on UK Article 17 reporting.

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

NATIONAL LEVEL

1. General information

1.1 Member State	UK
1.2 Species code	2016
1.3 Species scientific name	Pipistrellus kuhlii
1.4 Alternative species scientific name	
1.5 Common name (in national language)	Kuhl's pipistrelle

2. Maps

2.1 Sensitive species	No
2.2 Year or period	
2.3 Distribution map	No
2.4 Distribution map Method used	Insufficient or no data available
2.5 Additional maps	No

3. Information related to Annex V Species (Art. 14)

3.1 Is the species taken in the wild/exploited?	No	
3.2 Which of the measures in Art. 14 have been taken?	a) regulations regarding access to property	No
	b) temporary or local prohibition of the taking of specimens in the wild and exploitation	No
	c) regulation of the periods and/or methods of taking specimens	No
	d) application of hunting and fishing rules which take account of the conservation of such populations	No
	e) establishment of a system of licences for taking specimens or of quotas	No
	f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens	No
	g) breeding in captivity of animal species as well as artificial propagation of plant species	No
	h) other measures	No

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

b) Statistics/ quantity taken	Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period					
	Season/ year 1	Season/ year 2	Season/ year 3	Season/ year 4	Season/ year 5	Season/ year 6
Min. (raw, ie. not rounded)						
Max. (raw, ie. not rounded)						
Unknown	No	No	No	No	No	No

3.4. Hunting bag or quantity taken in the wild Method used

3.5. Additional information

BIOGEOGRAPHICAL LEVEL

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

Atlantic (ATL)

4.2 Sources of information

Dietz, C., Von Helverson, O & Nils, D. 2007. Handbuch der Fledermause Europas und Nordwestafrikas; Biologie, Kennzeichen, Gefährdung. Franckh-Kosmos, Stuttgart. 399 pp [Translation with extra text by Lina, P.H.C. & Hutson, A.M. 2009. Bats of Britain, Europe and North-West Africa. A&C Black, London. 400pp.]
 Dietz, C. & Kiefer, A. 2014. Die Fledermause Europas; Kennen, bestimmen, schützen. Franckh-Kosmos, Stuttgart. 394pp. [Translation with extra text by Lina, P.H.C. & Hutson, A.M. 2016. Bats of Britain and Europe. Bloomsbury, London. 398pp.]
 Harris, S. & Yalden, D.W. (Eds) 2008. Mammals of the British Isles: Handbook, 4th edition. The Mammal Society, Southampton. 799pp.
 Hutson, A.M. In prep. Scarce migrant, vagrant and imported bats in UK and Channel Islands.

5. Range

5.1 Surface area (km²)

5.2 Short-term trend Period

5.3 Short-term trend Direction

5.4 Short-term trend Magnitude

a) Minimum

b) Maximum

5.5 Short-term trend Method used

5.6 Long-term trend Period

5.7 Long-term trend Direction

5.8 Long-term trend Magnitude

a) Minimum

b) Maximum

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

5.9 Long-term trend Method used

5.10 Favourable reference range

- a) Area (km²)
- b) Operator
- c) Unknown
- d) Method

5.11 Change and reason for change in surface area of range

No change
The change is mainly due to:

5.12 Additional information

As this is a rare vagrant species, it is not appropriate to map this species' Range or set a value for the Favourable Reference Range.

6. Population

6.1 Year or period

6.2 Population size (in reporting unit)

- a) Unit
- b) Minimum
- c) Maximum
- d) Best single value

6.3 Type of estimate

6.4 Additional population size (using population unit other than reporting unit)

- a) Unit
- b) Minimum
- c) Maximum
- d) Best single value

6.5 Type of estimate

6.6 Population size Method used

6.7 Short-term trend Period

6.8 Short-term trend Direction

6.9 Short-term trend Magnitude

- a) Minimum
- b) Maximum
- c) Confidence interval

6.10 Short-term trend Method used

6.11 Long-term trend Period

6.12 Long-term trend Direction

6.13 Long-term trend Magnitude

- a) Minimum
- b) Maximum
- c) Confidence interval

6.14 Long-term trend Method used

6.15 Favourable reference population (using the unit in 6.2 or 6.4)

- a) Population size
- b) Operator
- c) Unknown
- d) Method

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

6.16 Change and reason for change in population size

No change
The change is mainly due to:

6.17 Additional information

As this is a rare vagrant species, it is not appropriate to estimate Population size and trend, or to set a value for the Favourable Reference Population.

7. Habitat for the species

7.1 Sufficiency of area and quality of occupied habitat

- a) Are area and quality of occupied habitat sufficient (for long-term survival)?
- b) Is there a sufficiently large area of unoccupied habitat of suitable quality (for long-term survival)?

7.2 Sufficiency of area and quality of occupied habitat Method used

7.3 Short-term trend Period

7.4 Short-term trend Direction

7.5 Short-term trend Method used

7.6 Long-term trend Period

7.7 Long-term trend Direction

7.8 Long-term trend Method used

7.9 Additional information

As this is a rare vagrant species, it is not appropriate to assess the Habitat for the species.

8. Main pressures and threats

8.1 Characterisation of pressures/threats

8.2 Sources of information

8.3 Additional information

9. Conservation measures

9.1 Status of measures

- a) Are measures needed? **No**
- b) Indicate the status of measures

9.2 Main purpose of the measures taken

9.3 Location of the measures taken

9.4 Response to the measures

9.5 List of main conservation measures

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

9.6 Additional information

10. Future prospects

- 10.1 Future prospects of parameters
- a) Range
 - b) Population
 - c) Habitat of the species

10.2 Additional information

As this is a rare vagrant species, it is not appropriate to consider the Future prospects.

11. Conclusions

11.1. Range

11.2. Population

11.3. Habitat for the species

11.4. Future prospects

11.5 Overall assessment of Conservation Status

11.6 Overall trend in Conservation Status

11.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

11.8 Additional information

As this is a rare vagrant species, an assessment of conservation status is not possible or necessary.

12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

- 12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)
- a) Unit
 - b) Minimum
 - c) Maximum
 - d) Best single value

12.2 Type of estimate

12.3 Population size inside the network Method used

12.4 Short-term trend of population size within the network Direction

12.5 Short-term trend of population size within the network Method used

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

12.6 Additional information

13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

A southern species that has undergone a rapid northerly expansion of range (particularly in the 1980s and 1990s) to include France (to Normandy) and the Channel Islands, and across to Ukraine. Summer roosts usually in buildings, winter roosts include rock fissures and cellars. Forages for small insects (mainly Diptera and Hymenoptera) over a variety of habitats, including water bodies, disturbed agriculture and urban habitats (and including around lights). Believed to be generally a more or less sedentary species. Since 1991, about 15 UK records, plus a number from Jersey (where it breeds) and Guernsey. Three or four of the UK records clearly imports, rest mainly south coast with one record from Cornwall, most around coast from Isle of Wight to London. A record from Felixstowe (Suffolk) may be an import. Apparent natural records mainly associated with migration or dispersal period. This species can currently be regarded as a rare migrant, but its potential to establish as a breeding species should not be overlooked.