

NORTH-WEST COAST OF LEWIS

J.E. Gordon

OS Grid Reference: NB512656

Highlights

The landforms and deposits on the coast of north-west Lewis provide important evidence for interpreting the patterns of sea-level change and glaciation in an area located towards the periphery of the main centres of glaciation. Part of the area remained ice-free throughout the Late Devensian.

Introduction

This site comprises a series of sections and landforms along a 16 km stretch of the north-west coast of Lewis, from Cunnadal (NB 512655) to Cladach Lag na Greine (NB 387557). These have long provided some of the most important evidence for interpreting the Pleistocene history of the Outer Hebrides (Geikie, 1873, 1874, 1877, 1878; Baden-Powell, 1938; McCann, 1968; von Weyarn, 1974, 1979; Peacock, 1984a, 1991; Sutherland and Walker, 1984). The interest comprises a fossil shore platform, pre-Holocene raised beach deposits, peat (?interglacial), till, complex glacial sediment sequences and solifluction deposits. Together, these features are of the very highest importance for studying glaciation and sea-level change in an area which has a fundamental bearing on the wider understanding of ice-movement patterns, ice sources, ice extent and climatic conditions in north-west Britain during the Pleistocene.

Description

Early descriptions recognized an essentially tripartite sequence in the area of Swainbost (NB 500635), comprising sands, gravels, clays and silts interbedded between two tills, with shells present in each of the layers (Geikie, 1873, 1874, 1877, 1878; Baden-Powell, 1938). However, more recent work has shown a much more complex succession of landforms and deposits represented at a number of localities along the coast (Figure 12.2). The principal features in the currently recognized sequence are as follows:




Figure 12.2: Quaternary deposits of north-west Lewis (from Peacock, 1984a).

5. till in the south and north; head and soliflucted till in the central part; multiple drift deposits at Swainbost.

4. raised beach deposits with a cryoturbated upper horizon

3. head

2. at different locations the shore platform is overlain by till and by peat; the relationship of the till to the peat is undetermined.

1. a raised shore platform and cliffline

The oldest marine feature clearly recognizable is a low-level (7–10 m O.D.) raised shore platform and cliffline (Godard, 1965; McCann, 1968; von Weymarn, 1974). It occurs discontinuously in north-west Lewis and on the Eye peninsula (Figure 12.1) but is best seen between Galson (NB 453603) and Dell (NB 472621) where it attains a width of 150 m (McCann, 1968; von Weymarn, 1974). As described by von Weymarn (1974), the principal features of the platform are:




Figure 12.1: Location map and principal Quaternary features of the Outer Hebrides (from Peacock, 1984a).

1. it is cut across the bedrock structure.

2. its seaward margin is sometimes covered in modern sand and shingle.

3. there is often a step down to a lower intertidal shore platform.

4. zones of weathered rock occur on its surface.

5. it is widest in embayments and narrows towards headlands where the backing cliff is best

seen. (eg. at the mouth of the Dibadall Dum at NB 470615 –McCann, 1968).

6. its landward margin is typically drift covered, although the backing cliff is locally seen (eg. at the mouth of the Dibadale Burn at NB 470615 –McCann, 1968).

7. abandoned stacks occur on its surface.

8. its distribution is largely confined to the area outside that of local glaciation in Lewis; a few remnants within the latter area are severely ice-modified.

Resting on the shore platform is a complex succession of deposits. In part, following von Weymarn (1974), these can be grouped into four units:

1. Raised beach deposits associated with till. Raised beach deposits form an important stratigraphic marker and were first identified at Galson where they appear to overlie till (Baden-Powell and Elton, 1937). Baden-Powell and Elton noted that Torridonian sandstone erratics were present in the beach deposits but absent from the till. The occurrence of the beach has since been identified intermittently along the whole coast (Figure 12.2) (McCann, 1968; von Weymarn, 1974; Peacock, 1984a). From the height of the one beach terrace of any extent at Galson, McCann (1968) inferred a maximum sea level of 32–37 ft (9.8–11.3 m) O.D. At several localities the upper layers of the beach deposit are cryoturbated. In addition to Galson, the raised beach deposits overlie till at Cladach na Luinge (NB 465611), Toa Dibadale (NB 469615), Cunndal and Swainbost (McCann, 1968; von Weymarn, 1974; Peacock, 1984a). In contrast, in the southern part of the site, to the south of Breivig (NB 414582), a separate till overlies the raised beach deposits.

2. Raised beach deposits associated with head and soliflucted till. Between North Galson (NB 438595) and Breivig, soliflucted till and head overlie the raised beach deposits; in other places the beach and soliflucted deposits are interbedded, for example at South Galson (NB 431591) and Cunndal.

3. At Toa Galson (NB 453603) an organic deposit interbedded with sand occurs above the shore platform and is overlain in turn by head and raised beach gravels (Figures 12.3 and 12.4). The organic deposit yielded radiocarbon dates greater than 39,100 to 47,150 BP (SRR–2365) (Sutherland and Walker, 1984). The pollen spectra (Figure 12.5) indicate development of the vegetation from treeless, open grassland to grassland with acid heath.




Figure 12.3: Toa Galson: sequence of sediments (from Sutherland and Walker, 1984).



Figure 12.4: Section at Toa Galson, north-west Lewis, showing the interglacial peat resting on bedrock and overlain by sand, head and the Galson Beach deposits. The upper part of the beach deposits has been affected by cryoturbation. (Photo: D. G. Sutherland.)




Figure 12.5: Relative pollen diagram for the peat deposit at Toa Galson showing selected taxa as percentages of total land pollen (from Sutherland and Walker, 1984). The location of the sample is shown in Figure 12.4.

4. Complex successions often referred to as multiple drift deposits. These occur in the Swainbost – Dell area and comprise interbedded till, sand, gravel, laminated clay and silt beds. Historically, they were described by Geikie (1873, 1874, 1877, 1878) and Baden-Powell (1938) who both recognized a tripartite sequence comprising sand, gravel, clay and silt interbedded between two till layers. Although more recent references acknowledge the complexity of the sequence (von Weymarn, 1974; Peacock, 1984a), there have been no detailed sedimentological studies. The deposits are glaciotectonically disturbed (Geikie, 1877; von Weymarn, 1974; Flinn, 1978b) and are rich in broken shell fragments. A faunal list published by Etheridge (1876) was updated by Baden-Powell (1938). According to Etheridge the faunal assemblage included species of arctic affinity (eg. *Astarte depressa* (Brown) and *Chlamys islandica* (Müller)) and northern affinity (eg. *Nuculana pernula* (Müller), *Arctica islandica* (L.), *Saxicava norvegica* Spengler, *Natica montacuti* Forbes and *Fusus gracilis* (Da Costa)); Baden-Powell inferred that only the upper till of the tripartite sequence contained a diagnostic faunal assemblage (eg. including *Astarte borealis* (Chemnitz), *Mya truncata* (L.) var. *uddevallensis* (Hancock), and *Panomya norvegica* (Spengler)), indicating cold water conditions. Radiocarbon dates on shells from the middle or lower layers of these deposits yielded ages of between 34,470 ± 720/-660 BP and 39,500 ± 1270/-1100 BP (inner fractions) (SRR-2368 to SRR-2370) (Sutherland and Walker, 1984).

Amino acid analyses of fragments of *Arctica islandica* from the same beds indicated that the shells were from at least two distinct periods. The older group (6 analyses) had alle/Ile ratios of 0.114 ± 0.007 and the younger group (9 analyses) had alle/Ile ratios of 0.079 ± 0.013 (D.G. Sutherland, unpublished data). Similar ratios on *Arctica islandica* from the last interglacial in the British Isles may be inferred to be 0.15–0.16 (Bowen and Sykes, 1988), indicating that the shells from north-west Lewis are Early and Middle Devensian.

Interpretation

Interpretations of the evidence in north-west Lewis fall into two main groups. The earlier accounts were focused on the drift sequences at Swainbost; later accounts have placed greater emphasis in the wider succession and its spatial variations.

In an early reference to the 'alluvial land' of north Lewis, McCulloch (1819) described superficial accumulations of 'clay and marle, together with a mixture of rolled stones of different kinds'. From their lithological composition he believed they were derived from the gneissic rocks of the mountains but was unable to suggest a mode of derivation.

Initially from analogies with Greenland glaciers Geikie (1874, p. 215) hinted that sequences such as the multiple drift deposits in north Lewis could have been laid down in a single episode at or near the front of a glacier terminating in the sea (see also Geikie, 1877, p. 185 footnote). Later, however, he interpreted the middle beds of the tripartite sequence as 'interglacial' marine deposits (Geikie, 1878) and postulated a succession of glacial and non-glacial episodes and relative sea-level fluctuations to explain the development of the full sequence over a much longer timescale (Geikie, 1877, 1878).

Geikie (1878) noted the restricted distribution of the shelly till and interbedded deposits. He believed that the latter had once been more extensive but had been removed by the ice which deposited the upper till; the shelly till was confined to the area where basal ice, deflected by the Hebridean land mass, impinged on its northern periphery.

Subsequently, Jehu and Craig (1934) and Baden-Powell (1938) acknowledged Geikie's interpretation of interglacial submergence. Baden-Powell suggested that the submergence was at least 17 m and possibly as much as 60 m, the latter value corresponding with one of the planation surfaces identified by Panzer (1928). Godard (1965) largely concurred, but raised the question as to whether the altitude of the marine deposits might simply reflect older marine material reworked by glacial meltwaters. More recent interpretations, however, are agreed that the multiple drift deposits of north-west Lewis are glacial in origin and were formed during a single glacial episode (von Weymarn, 1974; Flinn, 1978b; Peacock, 1984a; Sutherland and Walker, 1984).

Based on what was the first systematic account of the landforms and deposits in north-west

Lewis, von Weymarn (1974) inferred a possible sequence of events commencing with formation of the raised shore platform during one or more pre-Devensian interglacials. The platform was over-ridden by glacier ice from the east and buried by till containing Torridonian sandstone erratics. During the Ipswichian, relative sea level rose by around 7 m and the earlier till was reworked. During the Devensian glacial phases the raised beach gravels were over-ridden to the south of Borve by a local ice cap on Lewis, but the coast to the north remained free of ice and subject to intense periglacial activity. The multiple drift deposits probably related to external ice in the north of Lewis during the Late Devensian.

Alternative suggestions were made by Jardine (1977) that the beach deposits could date from the Loch Lomond Stadial rather than the Ipswichian, and by Dawson (1979c) that the solifluction deposits could be of Loch Lomond Stadial age and the beach formed earlier during the Lateglacial. Flinn (1978b) inferred that the multiple drift deposits at Swainbost were deposited prior to the last glacial maximum when the northern part of Lewis was covered by relatively inactive ice near the margin of the local ice cap. However, subsequent work by Peacock (1981a, 1984a) and Sutherland and Walker (1984) have substantially confirmed von Weymarn's interpretations and elaborated on the details. Peacock (1981a, 1984a) identified possible limits of the Late Devensian ice in north-west Lewis. The northern limit of the local ice cap was at Breivig. The limit of the external ice to the north was more difficult to define and he presented two alternative positions. The critical problem concerned the stratigraphic relationship of the raised beach to the multiple drift sequence at Swainbost. Peacock considered that the beach lay above the latter and therefore favoured a relatively small ice-covered area, leaving the coast between Breivig and the Butt of Lewis unglaciated. The alternative explanation that a lobe of ice crossed the coast in the Swainbost area was considered less probable because of the inferred stratigraphic relationship of the raised beach and multiple drift sequence.

Sutherland and Walker (1984) produced a different interpretation of the stratigraphy from that of Von Weymarn and Peacock, concluding that the multiple drift sequences on the north-west coast lay stratigraphically above the raised beach. A Late Devensian age for the multiple drift sequence is supported by the radiocarbon dates and amino acid analyses on the shells from the glacial deposits. Therefore the coast between Eoropie and Dell Sands, where the multiple drift sequences occur, was covered by the last ice sheet, confirming Peacock's alternative hypothesis.

The radiocarbon dates from the peat beneath the raised beach at Toa Galson confirm the antiquity of the deposit. Sutherland and Walker (1984) tentatively considered it to be of interglacial origin although there is no evidence to assign it to a particular stage. Significantly, the pollen spectra differ from those associated with the interstadial sites at Tolsta Head (see below) and St Kilda (Sutherland *et al.*, 1984), both of which produced finite radiocarbon dates.

North-west Lewis is therefore an area of the highest importance for demonstrating a series of glacial, periglacial and marine events of Devensian and possibly pre-Devensian ages. In sequence, the key elements are as follows:

6. Late Devensian features including till limits, shelly multiple drift sequences and ice-free areas with associated periglacial deposits.
5. Raised beach deposits (pre-Late Devensian).
4. Periglacial deposits.
3. Organic deposits, possibly interglacial (pre-dating the raised beach deposits).
2. Till (pre-dating the raised beach deposits).
1. Raised shore platform of pre-Devensian age.

Key localities within the site are as follows:

1. Cunndal, where till is overlain by raised beach which is succeeded by soliflucted till. Cunndal lies outside the Late Devensian ice margin.

2. Traigh Sands (NB 511644) to Peicir (NB 485625) which includes raised beach and the classic shelly multiple drift sequences now dated as Late Devensian in age.
3. Aird Dell to Toa Galson, which demonstrates the sequence of shore platform, till and raised beach outside the Late Devensian ice limit.
4. Immediately south of Toa Galson organic deposits occur between the shore platform and the raised beach.
5. Farther south, between North Galson and Breivig, the raised beach terrace is best developed and is overlain by soliflucted till and head.
6. At Breivig the Late Devensian ice limit crosses the coast and till overlies the raised beach here and to the south.
7. At Cladach Lag na Greine the raised beach rests on a shore platform and is overlain by till. Locally, where the platform rises above the beach deposit, it is striated, which provides important evidence that the beach was overridden by glacier ice and is overlain *in situ* lodgement, rather than soliflucted till.

The assemblage of deposits and landforms represented at these localities is unique in Scotland and has potential bearing on the interpretation of the Pleistocene history and chronology of events in north west Britain, including sea-level change, glacier reconstruction, palaeoclimatology and vegetation history. For example, as Sutherland and Walker (1984) point out, conventional models of the last Scottish ice sheet require re-assessment in the light of the evidence from north-west Lewis. The presence there of an ice-free area indicates a relatively restricted extent of the last ice sheet in Britain at its north-west margins, which contrasts with some recent reconstructions that placed the margin on, or near, the edge of the continental shelf (Boulton *et al.*, 1977; Andersen, 1981; Boulton *et al.*, 1985). It also supports a growing body of evidence (see Orkney, Caithness and north-east Scotland) that the northern margins of the last ice sheet may have been relatively limited in extent in contrast to the southern margins which reached south into South Wales, central England and the north coast of East Anglia. Sites such as north-west Lewis therefore provide critical field evidence to constrain and refine mathematical models of the last ice sheet and the palaeoclimatic inferences derived from them.

North-west Lewis is also unique in Scotland in providing unequivocal evidence for a Late Devensian ice-sheet limit on land and therefore provides exceptional opportunities for comparative studies of pre-Late Devensian and post-Late Devensian rates of weathering and soil development. Further, north-west Lewis is one of only a few sites in Scotland where interstadial or possibly interglacial organic deposits are represented. As such it is of the very highest importance for palaeoecological and palaeoenvironmental studies. Finally, the multiple drift deposits of north-west Lewis are of significant sedimentological interest for interpreting and reconstructing the depositional environments at the margin of the Late Devensian ice sheet.

Conclusions

The unique assemblage of landforms and deposits on the North-west Coast of Lewis has a long history of research and has provided critical information for interpreting the Pleistocene in Scotland. Among the key features of interest are a pre-Devensian shore platform, a pre-Late Devensian raised beach, organic deposits of possible interglacial origin, solifluction deposits and a complex sequence of fossiliferous sediments deposited by the Late Devensian ice-sheet (approximately 18,000 years ago). This area therefore provides a wealth of evidence for interpreting patterns of sea-level change, palaeoenvironmental conditions and ice-sheet history. One of the key findings to emerge from recent work is that part of the area remained ice-free when the Late Devensian glaciation was at its maximum extent (about 18,000 years ago).

Reference list

- Andersen, B.G. (1981) Late Weichselian ice sheets in Eurasia and Greenland. In *The Last Great Ice Sheets* (eds G.H. Denton and T.J. Hughes). New York, Wiley, 1–65.
- Baden-Powell, D.F.W. (1938) On the glacial and interglacial marine beds of north Lewis. *Geological Magazine*, **75**, 395–409.
- Baden-Powell, D.F.W. and Elton, C. (1937) On the relation between a raised beach and an Iron Age midden on the Island of Lewis, Outer Hebrides. *Proceedings of the Society of Antiquaries in Scotland*, **71**, 347–65.
- Boulton, G.S., Jones, A.S., Clayton, K.M. and Kenning, M.J. (1977) A British ice-sheet model and patterns of glacial erosion and deposition in Britain. In *British Quaternary Studies. Recent Advances* (ed. F.W. Shotton). Oxford, Clarendon Press, 231–46.
- Boulton, G.S., Smith, G.D., Jones, A.S. and Newsome, J. (1985) Glacial geology and glaciology of the last mid-latitude ice sheets. *Journal of the Geological Society*, **142**, 447–74.
- Bowen, D.Q. and Sykes, G.A. (1988) Correlation of marine events and glaciations on the north-east Atlantic margin. *Philosophical Transactions of the Royal Society of London*, **B318**, 619–35.
- Dawson, A.G. (1979c) Former sea-level changes in the Scottish Hebrides. *Hebridean Naturalist*, **3**, 16–22.
- Etheridge, R. (1876) Note on the fossils from the glacial deposits of the north-west coast of the Island of Lewis, Outer Hebrides. *Geological Magazine*, dec. 2, vol. 3, 552–5.
- Flinn, D. (1978b) The glaciation of the Outer Hebrides. *Geological Journal*, **13**, 195–9.
- Geikie, J. (1873) On the glacial phenomena of the Long Island or Outer Hebrides. First Paper. *Quarterly Journal of the Geological Society of London*, **29**, 532–45.
- Geikie, J. (1874) *The Great Ice Age and its Relation to the Antiquity of Man* 1st Edn. London, W. Isbister, 575pp.
- Geikie, J. (1877) *The Great Ice Age and its Relation to the Antiquity of Man* 2nd edn. London, Daldy, Isbister & Co. 624pp.
- Geikie, J. (1878) On the glacial phenomena of the Long Island or Outer Hebrides. Second Paper. *Quarterly Journal of the Geological Society of London*, **34**, 819–70.
- Godard, A. (1965) *Recherches de Géomorphologie en écosse du Nord-Ouest* Université de Strasburg, Publications de la Faculté des Lettres, Fondation Baulig, Tome 1, 701pp.
- Jardine, W.G. (1977) The Quaternary marine record in southwest Scotland and the Scottish Hebrides. In *The Quaternary of the Irish Sea* (eds C. Kidson and M.J. Tooley). Liverpool, Seel House Press, 99–118.
- Jehu, T.J. and Craig, R.M. (1934) Geology of the Outer Hebrides. Part V – North Harris and Lewis. *Transactions of the Royal Society of Edinburgh*, **57**, 839–74.
- McCann, S.B. (1968) Raised shore platforms in the Western Isles of Scotland. In *Geography at Aberystwyth. Essays Written on the Occasion of the Departmental Jubilee 1917–18 – 1967–68* (eds E.G. Bowen, H. Carter and J.A. Taylor). Cardiff, University of Wales Press, 22–34.
- Panzer, W. (1928) Zur Oberflächengestalt der Ausseren Hebriden, Beobachtungen und Fragen. *Zeitschrift für Geomorphologie*, **3**, 167–203.
- Peacock, J.D. (1981a) Report and excursion guide - Lewis and Harris. *Quaternary Newsletter*, **35**, 45–54.
- Peacock, J.D. (1984a) Quaternary geology of the Outer Hebrides. *Report of the British Geological Survey*, **16(2)**, 26p.
- Peacock, J.D. (1991) Glacial deposits of the Hebridean region. In *Glacial Deposits in Great Britain and Ireland* (eds J. Ehlers, P.L. Gibbard and J. Rose). Rotterdam, Balkema, 109–19.
- Sutherland, D.G. and Walker, M.J.C. (1984) A late Devensian ice-free area and possible interglacial site on the Isle of Lewis, Scotland. *Nature*, **309**, 701–3.
- Sutherland, D.G., Ballantyne, C.K. and Walker, M.J.C. (1984) Late Quaternary glaciation and environmental change on St Kilda, Scotland, and their palaeoclimatic significance. *Boreas*, **13**, 261–72.
- Von Weymarn, J.A. (1974) Coastline development in Lewis and Harris, Outer Hebrides, with particular reference to the effects of glaciation. Unpublished PhD thesis, University of Aberdeen.
- Von Weymarn, J. (1979) A new concept of glaciation in Lewis and Harris, Outer Hebrides. *Proceedings of the Royal Society of Edinburgh*, **B77**, 97–105.