

---

# MARYTON

*J.E. Gordon*

OS Grid Reference: NO683565

## Highlights

Deposits in the coastal section at Maryton include a sequence of estuarine sediments and buried peat. These deposits provide a record of the changes that occurred in sea level and coastal environmental conditions during the Holocene. The site is particularly notable for the evidence it displays for a major coastal flood in eastern Scotland during the middle Holocene.

## Introduction

The site at Maryton (NO 683565) is a cliff section in raised estuarine deposits on the west side of the Montrose Basin. It is important for demonstrating Holocene sea-level fluctuations in eastern Scotland and provides one of the best exposures showing the stratigraphic relationships of the estuarine (carse) deposits of the Main Postglacial Shoreline. It is also the only site known where a possible storm-surge or tsunami deposit dating to slightly prior to the maximum of the Main Postglacial Transgression is currently exposed. The site has been described by Smith *et al.* (1980) and discussed in a wider context by Smith *et al.* (1985a); it has also been illustrated in Smith *et al.* (1985a), Smith and Dawson (1990) and Long *et al.* (1989b).

## Description

The superficial deposits of the Montrose area were first described in detail by Howden (1868) and later summarised by him (Howden, 1886) in the following general succession:

6. Blown sand
5. Estuarine (carse) clay and sand
4. Peat
3. Glaciofluvial sands and gravels
2. Laminated marine clay
1. Till

The basal unit of till was overlain by laminated marine clay extending up the South Esk valley to an altitude of 40 ft (12.2 m). Although the contact between the two was nowhere exposed, Howden was in no doubt of their stratigraphic sequence. The clay (bed 2) contained abundant fossil remains of arctic molluscs which Howden (1868) listed (see Dryleys). Jamieson (1865) gave additional details, and Brady *et al.* (1874) listed the ostracods.

Howden described sands and gravels (bed 3) overlying the marine clay and the till, which extended as far inland as the mouths of the Highland glens. These gravels were part of the sequence first described by Buckland (1841a), and in mineralogical composition were identical to the moraines of the Highland glens. Howden suggested that the sands and gravels were formed by meltwaters at the end of the great ice-sheet glaciation and subsequently while valley glaciers remained in the glens. Cullingford and Smith (1980) have mapped these deposits to the west of the Montrose Basin and noted that four separate outwash terraces can be related to distinct shorelines at altitudes of between 15 m and 25 m O.D. around the Montrose Basin.

In a number of stream courses tributary to the South Esk, Howden observed peat (bed 4) resting on the marine clays and extending on to the gravels. At one locality the peat was overlain by carse clays. Generally, however, the latter deposits rested on the lower marine clay

and reached an altitude of 15 ft (4.6 m). Shells similar to those found in the present-day estuary were locally abundant in the carse clay. Details were given by Jamieson (1865) and Howden (1868). More recently, the pattern of Holocene sea-level changes in the Montrose area has been investigated in detail by Crofts (1971, 1972, 1974) and Smith and Cullingford (1985). In particular, Maryton has been described in detail by Smith *et al.* (1980). A sequence of marine and non-marine organic deposits infills a number of gullies dissecting Late Devensian marine and glaciofluvial terraces at the western end of the Montrose Basin (Cullingford and Smith, 1980; Smith *et al.*, 1980). A section in a terrace bluff at Maryton shows (Figure 14.5):



Figure 14.5: Section at Maryton. Laminated Late Devensian marine clays are overlain by peat then a layer of grey, micaceous, silty, fine sand interpreted as a tsunami deposit. Above the latter is silty peat then carse clay. (Photo: D. E. Smith.)

5.	Grey silty clay (carse)	0.85 m
4.	Peat	0.10 m
3.	Grey, micaceous, silty fine sand	0.18 m
2.	Peat	0.15 m
1.	Laminated, pink silty clay	0.20 m

The sequence in the section is broadly similar to that revealed in boreholes in the Fullerton area 0.5 km to the west (Smith *et al.*, 1980). Samples for pollen analysis were taken from the Maryton section and two coring sites at Fullerton by Smith *et al.* (1980). They recognized three pollen assemblage zones at Maryton (Figure 14.6). Zone M–1 includes poorly preserved pollen from most of the laminated, pink silty clay (bed 1). Although the pollen frequencies are low and the range of taxa restricted, open habitat conditions are indicated. Zone M–2 includes pollen from the horizons between the upper layers of the pink silty clay (bed 1) and the lowest part of the carse clay (bed 5). Contrasts in pollen frequency and assemblage suggest a break in deposition between Zones M–1 and M–2. The main characteristics of Zone M–2 are the consistent representation of *Betula* and *Pinus sylvestris*, variable amounts of *Corylus*-type and *Salix*-type pollen and the presence of *Ulmus* and *Quercus*. Several taxa also suggest the

proximity of damp conditions (Gramineae, Cyperaceae, Chenopodiaceae, Nymphaeaceae and *Filipendula*). Zone M–3 includes the greater part of the carse clay. The main representatives of M–3 are *Pinus sylvestris*, *Corylus*-type, *Betula*, *Ulmus* and *Quercus*. Other taxa (Nymphaeaceae and Cyperaceae) again indicate local damp conditions.




Figure 14.6: Maryton: relative pollen diagram showing selected taxa as percentages of total pollen (from Smith *et al.*, 1980).

Smith *et al.* (1980) correlated pollen zone M–1 with an equivalent zone from a similar, basal, laminated clay layer in one of the Fullerton boreholes. It was thought to reflect an open habitat, Late Devensian environment. Zones M–2 and M–3 also have equivalents with similar tree and herbaceous pollen in both the Fullerton boreholes. These assemblages indicate a boreal environment, corresponding with Holocene "chronozone" FI I of West (1970). An additional zone, corresponding with "chronozone" FI II, is represented in one of the Fullerton profiles.

Smith *et al.* (1980) obtained radiocarbon dates for several horizons at Maryton and Fullerton (Table 14.4).

## Interpretation

Sequences similar to those described by Howden (1868, 1886) from the Montrose area have long been recognized elsewhere in the Tay, Forth and Ythan estuaries. Together with the evidence from these other sites, the Montrose deposits have formed an important part of the wider interpretation of Lateglacial and Holocene sea-level changes in eastern Scotland (Jamieson, 1865; J. Geikie, 1874; Wright, 1937; Sissons, 1967a; Cullingford *et al.*, 1986, 1991). A full sequence of Lateglacial and Holocene deposits is nowhere exposed, but the section at Maryton demonstrates a key part of the succession spanning the period of the transgression associated with the Main Postglacial Shoreline; the Lateglacial part of the succession is represented at Dryleys (see above).

In the Maryton sequence the basal, laminated, silty clay is interpreted as a Late Devensian estuarine deposit (see Dryleys above) (Smith *et al.*, 1980). There is then a break in the sequence of deposits until peat began to form during the Holocene about 7340 BP. About 7140 BP, the peat accumulation was interrupted by deposition of the grey, micaceous, silty sand layer, before resuming until about 6980 BP when it was terminated by deposition of the carse.

On geomorphological and stratigraphic grounds Smith *et al.* (1980) correlated the carse surface in the Maryton area with the Main Postglacial Shoreline of eastern Scotland (see Sissons *et al.*, 1966). The radiocarbon dates show that the Main Postglacial Transgression associated with this shoreline was in progress at 7140 BP and possibly culminated between

about 6983 and 6704 BP. In the context of the age of the Main Postglacial Shoreline, these dates are older than those from stratigraphically similar sites in the Tay Estuary and eastern Fife (Chisholm, 1971; Cullingford *et al.*, 1980; Morrison *et al.*, 1981; Smith *et al.*, 1985b), but similar to the dates obtained in the Western Forth Valley (Sissons and Brooks, 1971).

The grey, micaceous, silty sand layer has recently attracted particular attention since it appears to represent a high-magnitude marine event, and Maryton has become a reference site for demonstrating its stratigraphic position and sedimentary characteristics. It has also been identified in boreholes at Fullerton, Dryleys and other sites around the margins of the Montrose Basin. At Puggieston (see Dryleys), the top and base of peats below and above the sand have been dated respectively to 7120 + 75 BP (SRR–2120) and 6850 + 75 BP (SRR–2119), dates that are indistinguishable from the similarly placed samples at Fullerton. A similar sand layer occupying the same stratigraphic position and giving approximately the same radiocarbon age has now been detected at a number of sites on the east coast of Scotland ranging from the Beaully Firth in the north to the Carse of Stirling in the south (see Western Forth Valley, Silver Moss and Barnyards). It has been argued that this unique deposit was possibly the result of a major storm surge in the North Sea Basin around 7000 BP (Smith *et al.*, 1985a; Haggart, 1988b) or, more probably, a tsunami associated with a major submarine slide on the Norwegian continental slope (Dawson *et al.*, 1988; Long *et al.*, 1989a, 1989b).

The section at Maryton therefore provides important stratigraphic evidence for Holocene sea-level changes in the Montrose Basin area. It is significant in several respects. First, as one of the best sections in eastern Scotland illustrating the Main Postglacial Transgression, Maryton is an integral member of a network of sites in this area (see Western Forth Valley, Pitlowie, Silver Moss, and Philorth Valley) which together have important research potential for establishing the diachroneity of the culmination of the Main Postglacial Transgression and the formation of the Main Postglacial Shoreline and therefore for elucidating the wider regional patterns of glacio-isostatic and eustatic changes. Second, since Maryton is the only location where the tsunami deposit is exposed, it is critically important in further studies of this event. Third, in a historical context, Maryton is also significant for its location in one of a number of key reference areas, recognized for over 100 years as illustrating the pattern of Lateglacial and Holocene sea-level change.

## Conclusions

Maryton forms part of a network of sites that is important for establishing the pattern of Holocene sea-level variations in eastern Scotland (that is, during the last 10,000 years). It is particularly notable for one of the best exposures in the estuarine deposits of the Main Postglacial Shoreline (an extensive raised shoreline that formed approximately 6800 years ago), and also the only available exposure of the deposits formed by a major flood that affected coastal areas around 7000 years ago.

## Reference list

- Brady, G.S., Crosskey, H.W. and Robertson, D. (1874) *A Monograph of the Post-Tertiary Entomostraca of Scotland*. London, Palaeontographical Society Monograph, 229 pp.
- Buckland, W. (1841a) On the evidences of glaciers in Scotland and the north of England. *Proceedings of the Geological Society of London*, **3**, 332–7.
- Chisholm, J.I. (1971) The stratigraphy of the post-glacial marine transgression in NE Fife. *Bulletin of the Geological Survey of Great Britain*, **37**, 91–107.
- Crofts, R.S. (1971) Coastal process and evolution around St Cyrus, Angus–Kincardine. Unpublished MLitt thesis, University of Aberdeen.
- Crofts, R.S. (1972) Coastal sediments and process around St Cyrus. In *North East Scotland Geographical Essays* (ed. C.M. Clapperton). Aberdeen, Department of Geography, University of Aberdeen, 15–9.
- Crofts, R.S. (1974) A method to determine shingle supply to the coast. *Transactions of the Institute of British Geographers*, **62**, 115–27.
- Cullingford, R.A. and Smith, D.E. (1980) Late Devensian raised shorelines in Angus and Kincardineshire, Scotland. *Boreas*, **9**, 21–38.

- Cullingford, R.A., Caseldine, C.J. and Gotts, P.E. (1980) Early Flandrian land and sea-level changes in lower Strathearn. *Nature*, **184**, 159–61.
- Cullingford, R.A. Firth, C.R. and Smith, D.E. (1986) *Relative Sea Level Changes in Eastern Scotland from the Loch Lomond Stadial to the Present A Summary of Present Knowledge*. Review Paper, IGCP Project 200 Field Meeting, Montrose, May 1986, 29pp.
- Cullingford, R.A. Smith, D.E. and Firth, C.R. (1991) The altitude and age of the Main Postglacial Shoreline in eastern Scotland. *Quaternary International*, **9**, 39–52.
- Dawson, A.G., Long, D. and Smith, D.E. (1988) The Storegga Slides: evidence from eastern Scotland of a possible tsunami. *Marine Geology*, **82**, 271–6.
- Geikie, J. (1874) *The Great Ice Age and its Relation to the Antiquity of Man* 1st Edn. London, W. Isbister, 575pp.
- Haggart, B.A. (1988b) The stratigraphy, depositional environment and dating of a possible tidal surge deposit in the Beaully Firth area, north-east Scotland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **66**, 215–30.
- Howden, J.C. (1868) On the superficial deposits at the estuary of the South Esk. *Transactions of the Edinburgh Geological Society*, **1**, 138–50.
- Howden, J.C. (1886) On the glacial deposits at Montrose. *Report of the British Association for the Advancement of Science for 1885*, 1040.
- Jamieson, T.F. (1865) On the history of the last geological changes in Scotland. *Quarterly Journal of the Geological Society of London*, **21**, 161–203.
- Long, D., Smith, D.E. and Dawson, A.G. (1989a) A Holocene tsunami deposit in eastern Scotland. *Journal of Quaternary Science*, **4**, 61–6.
- Long, D., Dawson, A.G. and Smith, D.E. (1989b) Tsunami risk in north-western Europe: a Holocene example. *Terra Nova*, **1**, 532–7.
- Morrison, J., Smith, D.E., Cullingford, R.A. and Jones, R.L. (1981) The culmination of the main postglacial transgression in the Firth of Tay area, Scotland. *Proceedings of the Geologists' Association*, **92**, 197–209.
- Sissons, J.B. (1967a) *The Evolution of Scotland's Scenery*. Edinburgh, Oliver and Boyd, 259pp.
- Sissons, J.B. and Brooks, C.L. (1971) Dating of early postglacial land and sea level changes in the western Forth Valley. *Nature Physical Science*, **234**, 124–7.
- Sissons, J.B., Smith, D.E. and Cullingford, R.A. (1966) Late-glacial and post-glacial shorelines in south-east Scotland. *Transactions of the Institute of British Geographers*, **39**, 9–18.
- Smith, D.E. and Cullingford, R.A. (1985) Flandrian relative sea-level changes in the Montrose Basin area. *Scottish Geographical Magazine*, **101**, 91–105.
- Smith, D.E., Morrison, J., Jones, R.L. and Cullingford, R.A. (1980) Dating the Main Postglacial Shoreline in the Montrose area, Scotland. In *Timescales in Geomorphology* (eds R.A. Cullingford, D.A. Davidson and J. Lewin). Chichester, John Wiley & Sons Ltd., 225–45.
- Smith, D.E., Cullingford, R.A. and Haggart, B.A. (1985a) A major coastal flood during the Holocene in eastern Scotland. *Eizeitalter und Gegenwart*, **35**, 109–18.
- Smith, D.E., Dawson, A.G., Cullingford, R.A. and Harkness, D.D. (1985b) The stratigraphy of Flandrian relative sea-level changes at a site in Tayside, Scotland. *Earth Surface Processes and Landforms*, **10**, 17–25.
- West, R.G. (1970) Pollen zones in the Pleistocene of Great Britain and their correlation. *New Phytologist*, **69**, 1179–83.
- Wright, W.B. (1937) *The Quaternary Ice Age*. 2nd edn. London, Macmillan and Co., 478pp.