
STRUIE CHANNELS

J. E. Gordon

OS Grid Reference: NH672789

Highlights

This site provides a good example of a meltwater channel system formed during the melting of the Late Devensian ice-sheet; such systems are relatively rare in northern Scotland.

Introduction

The site (NH 670790) is located in Strathrory on the south side of a col at c. 208 m OD between the Dornoch Firth to the north and the Cromarty Firth to the south. It provides a good example of a glacial meltwater channel system in the northern Highlands. The Struie Channels were first recorded by Peach *et al.* (1912) and subsequently have been mapped and described by J.S. Smith (1968) and Leftley (1991); otherwise they have attracted little published comment despite the relative scarcity of well-developed meltwater channel systems north of the Great Glen.

Description

The interest comprises a series of subparallel meltwater channels. The largest is up to 33 m deep, 89 m wide and 2.5 km long (Figure 7.10). In plan form, the channels show anastomosing and branching patterns, as well as parallel forms, and locally small cut-off loops lie perched above the main channel (Figure 7.10). According to J.S. Smith (1968), the channels originate at the lowest point of the col, but in fact they begin several hundred metres on the lee side and some 30–40 m above the lowest point. J.S. Smith (1968) also noted that another channel runs south from the next col to the west, and a cross channel extends eastwards from it to link with the Strathrory system. The channels are also associated with glaciofluvial deposits, including an esker at the south end of Loch Sheilan (NH 676780) and gravel terraces to the west in Strathrory, which have been partially quarried (Harris and Peacock, 1969; Mykura *et al.*, 1978), revealing glaciolacustrine deposits (Leftley, 1991).




Figure 7.10: Geomorphology of the Struie meltwater channels, Strathroy (from J.S. Smith, 1968; Leftley, 1991).

Interpretation

Peach *et al.* (1912) interpreted the Struie Channels as representing ice-marginal drainage at successive levels along the edge of a wasting glacier which occupied the valley of the Allt Dearg immediately to the north of Strathroy. However, on account of their apparent relationship to the col, J.S. Smith (1968) considered that the channels were of subglacial origin and were associated with his Fortrose stage of deglaciation, when they carried meltwater south from the Dornoch Firth to the Cromarty Firth across the low col into Strathroy and then towards Scotsburn (NH 720763). Leftley (1991) also interpreted the channels as subglacial in origin, formed during a late stage in the deglaciation of the last ice-sheet, when a lobe of ice extended across the col from the north into Strathroy; this occurred penecontemporaneously with the development of a series of ice-dammed lakes in Strathroy.

In their anastomosing forms and location on the lee side of a col, the Struie Channels are similar to the superimposed subglacial forms described by Clapperton (1968) from the Cheviots. Such channels are considered to reflect regional hydraulic gradients associated with active ice (*cf.* Sugden and John, 1976; Shreve, 1985a, 1985b). However, in other aspects, particularly their parallel forms, the Struie Channels resemble many of the channel systems described from lowland Scotland (Sissons, 1960, 1961a) and Scandinavia (Mannerfelt, 1945, 1949), which have been interpreted as marginal or submarginal in origin. The Struie site therefore provides an interesting assemblage of meltwater channel features that would benefit from further detailed investigation.

Meltwater channel systems are relatively rare in the northern Highlands of Scotland, but Struie is a particularly good example. It demonstrates many of the typical features of meltwater

channels in Scotland (see Carlops and Rammer Cleugh), including a combination of subglacial and marginal/submarginal characteristics. In the wider context of the Moray Firth area, the Struie Channels complement the interests of the depositional, glaciofluvial landform assemblages at Torvean, Kildrummie Kames and Littlemill.

Conclusions

The Struie channels were eroded by glacial rivers during the melting of the last (Late Devensian) ice-sheet, between approximately 14,000 and 13,000 years ago. They form part of a network of sites showing glacial meltwater landforms formed at this time, and are notable as one of the few well-known examples of a system of meltwater channels in northern Scotland.

Reference list

- Clapperton, C.M. (1968) Channels formed by the superimposition of glacial meltwater streams with special reference to the east Cheviot Hills, north-east England. *Geografiska Annaler*, **50A**, 207–20.
- Harris, A.L. and Peacock, J.D. (1969) Sand and gravel resources of the inner Moray Firth. *Report of the Institute of Geological Sciences* **69/9**, 18pp.
- Leftley, D.C. (1991) The Late Devensian development of the Strathroy River Valley. Unpublished BSc dissertation, West London Institute of Higher Education.
- Mannerfelt, C.M. (1945) Några glacialmorfologiska formelement. *Geografiska Annaler*, **27**, 1–239.
- Mannerfelt, C.M. (1949) Marginal drainage channels as indicators of the gradients of Quaternary ice-caps. *Geografiska Annaler*, **31**, 194–9.
- Mykura, W., Ross, D.L. and May, F. (1978) Sand and gravel resources of the Highland Region. *Report of the Institute of Geological Sciences* **78/8**, 60pp.
- Peach, B.N., Gunn, W., Clough, C.T., Hinxman, L.W., Crampton, C.B. and Anderson, E.M. (1912) The geology of Ben Wyvis, Carn Chuinneag, Inchbae and the surrounding country, including Garve, Evanton, Alness and Kincardine. (Explanation of Sheet 93). Memoirs of the Geological Survey of Scotland. Edinburgh, HMSO, 189pp.
- Shreve, R.L. (1985a) Esker characteristics in terms of glacier physics, Katahdin esker system, Maine. *Geological Society of America Bulletin*, **96**, 639–46.
- Shreve, R.L. (1985b) Late Wisconsin ice-surface profile calculated from esker paths and types, Katahdin Esker System, Maine. *Quaternary Research*, **23**, 27–37.
- Sissons, J.B. (1960) Some aspects of glacial drainage channels in Britain. Part I. *Scottish Geographical Magazine*, **76**, 131–46.
- Sissons, J.B. (1961a) Some aspects of glacial drainage channels in Britain. Part II. *Scottish Geographical Magazine*, **77**, 15–36.
- Smith, D.E. (1968) Post-glacial displaced shorelines in the surface of the carse clay on the north bank of the River Forth, in Scotland. *Zeitschrift für Geomorphologie*, NF, **12**, 388–408.
- Sugden, D.E. and John, B.S. (1976) *Glaciers and landscape*. London, Edward Arnold, 376pp.