

# LOCH AN T-SUIDHE

M.J.C. Walker

OS Grid Reference: NM370215

## Highlights

The sediments that infill the floor of this loch contain a valuable pollen record, supported by radiocarbon dating, of vegetational history and environmental change on Mull during the Lateglacial.

## Introduction

Loch an t-Suidhe (NM 370215) is a small lochan at an altitude of 30 m OD, approximately 1 km west-south-west of Bunesan on the Ross of Mull. Although a large number of pollen sites have now been investigated on the Scottish mainland (Walker, 1984b), until recently relatively little information was available about the Lateglacial on the islands of the Inner and Outer Hebrides. The publication (Lowe and Walker, 1986a) of two radiocarbon-dated pollen diagrams from sites on the Isle of Mull was therefore of significance in the context of the Scottish Lateglacial. Of the two sites investigated, Loch an t-Suidhe offered the better stratigraphic resolution and a coherent series of six radiocarbon dates was obtained from the profile (Walker and Lowe, 1982). The wealth of palaeoenvironmental evidence contained within the sediments of this basin make it one of the most important Lateglacial pollen sites so far described from western Scotland.

## Description

The sedimentary sequence (Figure 11.15) near the southern shore of the lochan clearly resembles the tripartite lithostratigraphic Lateglacial succession commonly found in infilled Lateglacial lake sites in Britain (Sissons *et al.*, 1973). A lower minerogenic unit (approximately 8.8 m to base) is overlain by organic sediments (8.2–8.8 m) and these, in turn, are succeeded by a further minerogenic suite (7.9–8.2 m). The whole sequence is overlain by Holocene lake muds and peats. Details of the pollen stratigraphy of the Holocene sediments are contained in Lowe and Walker (1986b).




Figure 11.15: Loch an t-Suidhe: relative pollen diagram showing selected taxa as percentages of total land pollen (from Lowe and Walker, 1986a).

The pollen diagram from the Lateglacial and early Holocene deposits was divided into a series

of local pollen assemblage zones based on fluctuations in the curves for the principal taxa (Figure 11.15). These show (from the base) a sequence of increasing pollen content and diversity accompanied by rising curves for woody plant pollen (zones LSa to LSd; the Lateglacial Interstadial), followed by a phase of reduced woody plant pollen and an increase in pollen from taxa indicative of bare or disturbed soils (zones LSe and LSf; the Loch Lomond Stadial). This, in turn, is followed by a series of pollen assemblage zones dominated by successive maxima in *Empetrum*, *Juniperus*, *Betula* and *Corylus* (zones LSg to LSj; the early Holocene).

## Interpretation

In terms of regional landscape changes, the pollen record from Loch an t-Suidhe reflects initial vegetational colonization of freshly exposed substrates during the early Lateglacial Interstadial following the wastage of the Late Devensian ice-sheet. Subsequently, there developed a juniper scrub and *Empetrum* heath vegetation cover. This was succeeded by a marked vegetational 'revertance' phase (during pollen zones LSe and LSf) in response to the harsh climatic conditions of the Loch Lomond Stadial, when a tundra landscape developed. Finally, woody plants re-expanded as climate improved at the close of the Lateglacial and there occurred a vegetational succession from open heathland to *Betula*–*Corylus* woodland during the early Holocene.

Although this sequence is very similar to that recorded at Mishnish in northern Mull (Lowe and Walker, 1986a) and is comparable in broad outline with many other Lateglacial pollen successions from northern Britain (Pennington *et al.*, 1972; Gray and Lowe, 1977b; Pennington, 1977a, 1977b; Walker and Lowe, 1990), a number of features combine to make the Lateglacial and early Holocene record from Loch an t-Suidhe particularly distinctive. First, the data on deteriorated pollen provide independent evidence of episodes of increased geomorphological activity around the basin (for example, accelerated minerogenic inwash during the Loch Lomond Stadial). In addition, variations in levels of deterioration between individual pollen taxa made possible the differentiation between primary and secondary components in the pollen spectra, an aspect of pollen analysis that is becoming increasingly important in understanding the plant communities that developed during the Loch Lomond Stadial (Walker and Lowe, 1990). Second, following the work of Mackereth (1965, 1966) and Pennington *et al.* (1972), fluctuations in the curves for organic carbon and for the chemical elements Na, K, Mg and Ca, provide further evidence of the extent of mineral inwash into the basin and hence constitute an additional indirect record for landscape change around the site. Third, the fall in *Juniperus* pollen during the mid-interstadial and its replacement by *Empetrum*, in association with other evidence for disturbance in the vegetation cover, appears to reflect climatic deterioration some 1000 years before the onset of the Loch Lomond Stadial. This inference is in broad agreement with coleopteran records, which show evidence of a cooling trend, and particularly a fall in winter temperatures, from c. 12,500 BP onwards (Atkinson *et al.*, 1987). Fourth, systematic analysis of the record of *Artemisia* pollen reveals a change from oceanic conditions to a more continental climatic regime during the course of the Loch Lomond Stadial, which appears to be related to the southward migration of the oceanic and atmospheric polar fronts (Duplessy *et al.*, 1981; Ruddiman and McIntyre, 1981b; Bard *et al.*, 1987).

Finally, Loch an t-Suidhe is notable because of the internally consistent series of radiocarbon dates that was obtained from the profile. As with all age determinations on bulk samples of organic lake muds, however, contamination by older or younger carbon residues (Sutherland, 1980; Walker and Harkness, 1990) cannot be entirely excluded, and hence the dates must be treated with a degree of caution. The basal organic sediments were dated at 13,140 ± 100 BP (SRR-1805), an age determination which is in broad agreement with dates on comparable biostratigraphic horizons at sites on the Scottish mainland (Bishop, 1963; Kirk and Godwin, 1963; Sissons and Walker, 1974; Pennington, 1975b; Lowe and Walker, 1977; Vasari, 1977). It is also in agreement with the date inferred for the replacement of polar by warmer waters around the shores of western Britain (Duplessy *et al.*, 1981; Ruddiman and McIntyre, 1981b; Peacock and Harkness, 1990). If correct, the date from the basal sediments would support the view that climatic amelioration at the beginning of the Lateglacial Interstadial occurred around 13,000 BP (Coope, 1975; Atkinson *et al.*, 1987). The *Juniperus* decline in the interstadial was dated at 11,860 ± 80 BP (SRR-1804), an age determination that accords with the inference

that climatic deterioration in western Britain began around 12,000 BP (Watts, 1977, 1985; Craig, 1978; Walker and Lowe, 1990). The onset of the Loch Lomond Stadial (as indicated by the sediment and pollen records) was dated at 10,690 + 70 BP (SRR-1803), which is very close to the age determination (10,730 + 60 BP; SRR-1807) on the comparable horizon at Mishnish in northern Mull (Walker and Lowe, 1982). Three dates of 10,440 + 80 BP (SRR-1802), 10,200 + 70 BP (SRR-1801) and 10,200 + 70 BP (SRR-1800) were obtained from the early Holocene sediments which pre-date the expansion of *Juniperus*. These dates are in broad agreement with a number of dates from basal Holocene sediments from other Scottish sites (Walker and Lowe, 1979, 1980, 1985) and, if correct, they reinforce the suggestion (Lowe and Walker, 1976; Walker and Lowe, 1981) that climatic amelioration at the close of the Loch Lomond Stadial occurred well before 10,000 BP. It should be noted, however, that the recently discovered plateau of constant radiocarbon age around 10,000 BP, and which appears to reflect fluctuations in atmospheric radiocarbon production (Ammann and Lotter, 1989; Zbinden *et al.*, 1989), poses a major difficulty in the establishment of 'reliable' age estimates at the Lateglacial–Holocene boundary (see also Kingshouse).

Loch an t-Suidhe contains a wealth of data on the Lateglacial and early Holocene environmental history of the Isle of Mull. It forms a key element in a network of published sites from the mainland (Pennington *et al.*, 1972; Pennington, 1975b, 1977a; Rymer, 1977; Tipping, 1984) and from the Isle of Skye (Birks, 1973; Walker and Lowe, 1990) which now enable a regional picture to be established of environmental change along the western Scottish seaboard following the wastage of the last ice-sheet. However, Loch an t-Suidhe may also be important in a wider context. Although it is now generally recognized that a dominant influence on the British climate and environment has been oceanographic changes in the north-east Atlantic province (see, for example, Lowe and Walker, 1984), it has not always proved possible to establish clear links between the marine and terrestrial records, particularly over the relatively restricted timespan of the Lateglacial. The location of Loch an t-Suidhe on the maritime fringes of north-west Britain, coupled with the detailed stratigraphic evidence contained within the profile and the internally consistent radiocarbon chronology, make it a potentially valuable site for correlation between the marine and terrestrial records. In this respect, therefore, Loch an t-Suidhe may prove to be a site of both national and international significance.

## Conclusions

Loch an t-Suidhe is a key site for interpreting Lateglacial environmental history in western Scotland during the Lateglacial, between about 13,000 and 10,000 years ago. The pollen record shows the pattern of vegetation colonization and development in the period after melting of the last ice-sheet (about 13,000 years ago). There is clear evidence for a subsequent return to tundra conditions during the Loch Lomond Stadial (approximately 11,000 and 10,000 years ago), followed by the development of open heathland and birch woodland during the early Holocene. Loch an t-Suidhe is particularly significant for the wealth of information it provides about this important time period, allowing detailed reconstruction of the palaeoenvironmental conditions.

## Reference list

- Ammann, B. and Lotter, A.F. (1989) Late-Glacial radiocarbon and palynostratigraphy on the Swiss plateau. *Boreas*, **18**, 109–26.
- Atkinson, T.C., Briffa, K.R. and Coope, G.R. (1987) Seasonal temperatures in Britain during the last 22,000 years, reconstructed using beetle remains. *Nature*, **325**, 587–93.
- Bard, E., Arnold, M., Maurice, P., Duprat, J., Moyes, J. and Duplessy, J.-C. (1987) Retreat velocity of the North Atlantic polar front during the last deglaciation determined by  $^{14}\text{C}$  accelerator mass spectrometry. *Nature*, **328**, 791–4.
- Birks, H.J.B. (1973) *Past and Present Vegetation of the Isle of Skye. A Palaeoecological Study*. Cambridge, Cambridge University Press, 415pp.
- Bishop, W.W. (1963) Lateglacial deposits near Lockerbie, Dumfriesshire. *Transactions of the Dumfriesshire and Galloway Natural History and Antiquarian Society*, **40**, 117–32.
- Coope, G.R. (1975) Climatic fluctuations in northwest Europe since the last interglacial, indicated by fossil assemblages of coleoptera. In *Ice Ages: Ancient and Modern* (eds A.E.

- Wright and F. Moseley). Liverpool, Seel House Press, 153–68.
- Craig, A.J. (1978) Pollen percentage and influx analyses in south-east Ireland: a contribution to the ecological history of the Late-glacial period. *Journal of Ecology*, **66**, 297–324.
- Duplessy, J.-C., Delibrias, G., Turon, J.L., Pujol, C., and Duprat, J. (1981). Deglacial warming of the north-eastern Atlantic Ocean; correlation with the palaeoclimatic evolution of the European continent. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **35**, 121–44.
- Gray, J.M. and Lowe, J.J. (1977b) The Scottish Lateglacial environment: a synthesis. In *Studies in the Scottish Lateglacial Environment* (eds J.M. Gray and J.J. Lowe). Oxford, Pergamon Press, 163–81.
- Kirk, W. and Godwin, H. (1963) A late glacial site at Loch Droma, Ross and Cromarty. *Transactions of the Royal Society of Edinburgh*, **65**, 225–49.
- Lowe, J.J. and Walker, M.J.C. (1976) Radiocarbon dates and deglaciation of Rannoch Moor, Scotland. *Nature*, **246**, 632–3.
- Lowe, J.J. and Walker, M.J.C. (1977) The reconstruction of the Lateglacial environment in the southern and eastern Grampian Highlands. In *Studies in the Scottish Lateglacial Environment* (eds J.M. Gray and J.J. Lowe). Oxford, Pergamon Press, 101–18.
- Lowe, J.J. and Walker, M.J.C. (1984) *Reconstructing Quaternary Environments*. London, Longman, 389pp.
- Lowe, J.J. and Walker, M.J.C. (1986a) Lateglacial and early Flandrian environmental history of the Isle of Mull, Inner Hebrides, Scotland. *Transactions of the Royal Society of Edinburgh Earth Sciences*, **77**, 1–20.
- Lowe, J.J. and Walker, M.J.C. (1986b) Flandrian environmental history of the Isle of Mull, Scotland. II. Pollen analytical data from sites in western and northern Mull. *New Phytologist*, **103**, 417–36.
- Mackereth, F.J. (1965) Chemical investigations of lake sediments and their interpretation. *Proceedings of the Royal Society London*, **161**, 295–309.
- Mackereth, F.J. (1966) Some chemical observations on post-glacial lake sediments. *Philosophical Transactions of the Royal Society of London* **B250**, 165–213.
- Peacock, J.D. and Harkness, D.D. (1990) Radiocarbon ages and the full-glacial to Holocene transition in seas adjacent to Scotland and southern Scandinavia: a review. *Transactions of the Royal Society of Edinburgh. Earth Sciences*, **81**, 385–96.
- Pennington, W. (1975b) A chronostratigraphic comparison of Late-Weichselian and Late-Devensian subdivisions, illustrated by two radiocarbon-dated profiles from western Britain. *Boreas*, **4**, 157–71.
- Pennington, W. (1977a) Lake sediments and the Lateglacial environment in northern Scotland. In *Studies in the Scottish Lateglacial Environment* (eds J.M. Gray and J.J. Lowe). Oxford, Pergamon Press, 119–41.
- Pennington, W. (1977b) The Late Devensian flora and vegetation of Britain. *Philosophical Transactions of the Royal Society*, **280B**, 247–71.
- Pennington, W., Haworth, E.Y., Bonny, A.P. and Lishman, J.P. (1972) Lake sediments in northern Scotland. *Philosophical Transactions of the Royal Society of London* **264B**, 191–294.
- Ruddiman, W.F. and McIntyre, A. (1981b) The North Atlantic during the last deglaciation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **35**, 145–214.
- Rymer, L. (1977) A late-glacial and early post-glacial pollen diagram from Drimnagall, North Knapdale, Argyllshire. *New Phytologist*, **79**, 211–21.
- Sissons, J.B. and Walker, M.J.C. (1974) Lateglacial site in the central Grampian Highlands. *Nature*, **249**, 822–4.
- Sissons, J.B., Lowe, J.J., Thompson, K.S.R. and Walker, M.J.C. (1973) Loch Lomond Readvance in the Grampian Highlands of Scotland. *Nature Physical Science*, **244**, 75–7.
- Sutherland, D.G. (1980) Problems of radiocarbon dating deposits from newly deglaciated terrain: examples from the Scottish Lateglacial. In *Studies in the Lateglacial of Northwest Europe* (eds J.J. Lowe, J.M. Gray and J.E. Robinson). Oxford, Pergamon Press, 139–49.
- Tipping, R.M. (1984) Late Devensian and Early Flandrian vegetational history and deglacial chronology of western Argyll. Unpublished PhD thesis, City of London Polytechnic.
- Vasari, Y. (1977) Radiocarbon dating of the Lateglacial and early Flandrian vegetational succession in the Scottish Highlands and the Isle of Skye. In *Studies in the Scottish Lateglacial Environment* (eds J.M. Gray and J.J. Lowe). Oxford, Pergamon Press, 143–62.
- Walker, M.J.C. (1984b) Pollen analysis and Quaternary research in Scotland. *Quaternary Science Reviews*, **4**, 369–404.

- 
- Walker, M.J.C. and Harkness, D.D. (1990) Radiocarbon dating the Devensian Lateglacial in Britain: new evidence from Llanilid, South Wales. *Journal of Quaternary Science*, **5**, 135–44.
- Walker, M.J.C. and Lowe, J.J. (1979) Postglacial environmental history of Rannoch Moor, Scotland. II. Pollen diagrams and radiocarbon dates from the Rannoch Station and Corroun areas. *Journal of Biogeography*, **6**, 349–62.
- Walker, M.J.C. and Lowe, J.J. (1980) Pollen analyses, radiocarbon dates and the deglaciation of Rannoch Moor, Scotland, following the Loch Lomond Advance. In *Timescales in Geomorphology* (eds R.A. Cullingford, D.A. Davidson and J. Lewin). Chichester, John Wiley, 247–59.
- Walker, M.J.C. and Lowe, J.J. (1981) Postglacial environmental history of Rannoch Moor, Scotland. III. Early- and mid-Flandrian pollen stratigraphic data from sites on western Rannoch Moor and near Fort William. *Journal of Biogeography*, **8**, 475–91.
- Walker, M.J.C. and Lowe, J.J. (1982) Lateglacial and early Flandrian chronology of the Isle of Mull, Scotland. *Nature*, **296**, 558–61.
- Walker, M.J.C. and Lowe, J.J. (1985) Flandrian environmental history of the Isle of Mull, Scotland. I. Pollen-stratigraphic evidence and radiocarbon dates from Glen More, south-central Mull. *New Phytologist*, **99**, 587–610.
- Walker, M.J.C. and Lowe, J.J. (1990) Reconstruction of the environmental history of the last glacial-interglacial transition: evidence from the Isle of Skye, Inner Hebrides, Scotland. *Quaternary Science Reviews*, **9**, 15–49.
- Watts, W.A. (1977) The Late Devensian vegetation of Ireland. *Philosophical Transactions of the Royal Society of London*, **B280**, 273–93.
- Watts, W.A. (1985) Quaternary vegetation cycles. In *The Quaternary History of Ireland* (eds K.J. Edwards and W.P. Warren). London, Academic Press, 155–85.
- Zbinden, H., Andrée, M., Oeschger, H., Ammann, B., Lotter, A., Bonani, G. and Wölfli, W. (1989) Atmospheric radiocarbon at the end of the last glacial: an estimate based on AMS radiocarbon dates on terrestrial macrofossils from lake sediments. *Radiocarbon*, **31**, 795–804.