

---

# AGASSIZ ROCK

*J.E. Gordon*

OS Grid Reference: NT259702

## Highlights

The striated rock surface at Agassiz Rock was first recognised by Louis Agassiz in 1840 to have been eroded by glacier ice. The site is historically significant for its part in the development of the glacial theory in Scotland.

## Introduction

Agassiz Rock (NT 254702) is located on the south side of Blackford Hill in Edinburgh. It is principally of historical interest for a striated rock surface that was associated with the early development of the glacial theory in Scotland. In addition, Agassiz Rock represents an important landmark in geological conservation, being one of the earliest Quaternary sites recognised to require safeguarding. The site has been referred to in a number of papers (Rhind, 1836; Milne Home, 1840, 1846, 1847a) Maclaren, 1841, 1842a; Buckland, 1841a; Fleming, 1859; Panton, 1873; Brown, 1874; Peach *et al.*, 1910; Mitchell and Mykura, 1962). It was included in field excursions of the 1948 International Geological Congress in Britain (International Geological Congress, 1948) and, in addition, it features in the itineraries recommended by Geikie (1901), Campbell (1951) and Waterston (1960).

## Description

Agassiz Rock is located on the south side of Blackford Hill where an andesite cliff has been undercut to form a shallow cave, the rock surfaces of which are grooved and striated like the overhanging cliff (Figure 17.2). Early descriptions of the site include those of Rhind (1836) and Milne Home (1840). The former explained the grooving by molten rock falling on a bed of sand and retaining the moulded impression of its surface; the latter in terms of marine submergence. However, it was on 27 October, 1840, that the site attained its fame when it was visited by Louis Agassiz. A few weeks earlier, Agassiz had delivered a paper at a British Association meeting in Glasgow in which he argued that all the northern parts of Europe, Asia and America were formerly covered with a mass of ice (Agassiz, 1841a). Although it is not recorded in the abstract of his paper, Agassiz apparently alluded to the former existence of glaciers in Scotland (Anon, 1840; Maclaren, 1840). After the meeting he departed on a tour of Scotland to investigate the field evidence (Davies, 1968a, 1968b). In the course of this journey, accompanied by William Buckland, he found many striking and convincing traces of former glaciers. When he visited Edinburgh he was taken on a tour to search for glacier markings on the south side of the city by a group of Edinburgh geologists, including Charles Maclaren, then editor of *The Scotsman*. Agassiz was doubtful about some of the features initially shown to him, but on seeing the cave at Blackford Hill is reputed to have exclaimed "That is the work of the ice" (Maclaren, 1841, 1842a; Cox and Nicol, 1869).



Figure 17.2: Part of the smoothed and grooved rock surface at Agassiz Rock, Edinburgh, which has been attributed to glacial abrasion. The form of the rock surface bears a strong resemblance to glacially abraded surfaces elsewhere in Scotland and in modern glacial environments. (Photo: J. E. Gordon.)

The striations at Agassiz Rock from part of a local assemblage of features that indicate ice moving eastwards across the area (Figure 17.3). Blackford Hill itself is a crag and tail, 1.5 km long, with deep erosional grooves on both its north and south sides, comparable to those around Edinburgh Castle Rock (Sissons, 1971). A smaller superimposed crag and tail occurs at Corbie's Craig south of the hill top, and to the east, clast fabric measurements in the main drift tail also conform with ice flowing to the east (Kirby, 1969b).




Figure 17.3: Blackford Hill crag and tail, showing a till clast fabric in the tail, the Corbie's Craig crag and tail and the direction of ice movement inferred from the striations at Agassiz Rock (from Kirby, 1969b).

In the Edinburgh area, a prominent theme in many of the earlier 19th century accounts is the recognition of the overall easterly movement of the agent responsible for the superficial deposits and bedrock striations (Figure 17.4). Typical lines of evidence included the disposition

of crag-and-tail forms (Hall, 1815; Maclaren, 1828, 1866); the transport of erratics (Milne Home, 1840, 1871, 1874a, 1874b; Nicol, 1848; Fleming, 1859; Campbell and Anderson, 1909); deformation and overfolding of strata to the east (Milne Home, 1840, 1871; Fleming, 1859; Brown, 1874); bedrock striations and moulding (Imrie, 1812; Hall, 1815; Maclaren, 1828, 1842b, 1866; Milne Home, 1840; Fleming, 1847, 1859; Chambers, 1853; Miller, 1864; Henderson, 1872; Richardson, 1877a, 1877b; Goodchild, 1896); and striations on stones in till (Maclaren, 1849; 1866; Milne Home, 1840; Henderson, 1874; Miller, 1864; Miller, 1884). Miller (1884) produced the first map showing the pattern of striations on bedrock and till clasts. Further instances of striations and movements of erratic are given by Peach *et al.* (1910a), and Peach (1909) described the classic Lennoxton essexite boulder train (see also Shakesby 1978, 1979, 1981). More recently Burke (1968, 1969) has quantified some of the evidence for these ice movement trends, and Sissons (1971) has described the strong imprint of glacial erosion in central Edinburgh.




Figure 17.4: Indicators of ice movement in the Lothians area recorded up until 1863 (from Kirby, 1969b).

## Interpretation

Although Agassiz Rock does not bear the distinction of being the first site in Scotland to have been recognised as the product of land ice, it was nevertheless of considerable significance (Buckland, 1841a; Davies, 1968a, 1968b), since the striations under the overhang could not have been produced by marine-drifted icebergs, the hypothesis of many contemporary geologists to explain such phenomena. Nor could they have been formed by debris-laden catastrophic deluges or floods as suggested by Hall (1815) to explain striations on nearby Corstorphine Hill because of their close parallel arrangement over short distances.

Nevertheless, Milne Home (1846, 1847a) and Fleming (1859) were not convinced of the glacial origin of the striations at Blackford Hill and other localities around Edinburgh. The former persisted with the diluvial hypothesis, and the latter explained them as a local phenomenon associated with the Braid Burn. Geikie (1863a), however, in his important exposition on the evidence for former glaciers in Scotland clearly established that striations, including those at Blackford Hill, were the product of land ice. Brown (1874) also believed the striations to be glacial but considered that a large landslip had brought the striated rock to its present position.

Subsequent references in the literature to Agassiz Rock (Panton, 1873; Peach *et al.*, 1910a;

Mitchell and Mykura, 1962) acknowledge the historical significance of the site, although critics have suggested that some of the striations may in fact be tectonic slickensides (Mitchell and Mykura, 1962).

Agassiz Rock is a site of considerable historical interest as one of the classic localities that played a significant part in the development of the glacial theory in Scotland. Its striated rock surface was among the first such features to be recognised as the product of glacier ice by Louis Agassiz in 1840. It is also significant in another historical context, being one of the first geological sites recognised to require practical measures to ensure the preservation of its interest. In 1908 the Council of Edinburgh Geological Society successfully negotiated with Edinburgh Town Council to place a railing around the site and erect a memorial tablet (Watson, 1934). The railing is now dismantled, and a new plaque erected by the Nature Conservancy Council reads "Agassiz Rock. In 1840 Louis Agassiz Swiss Geologist stated that this rock was polished and grooved by ice during the great ice age".

## Conclusions

Agassiz Rock is a site of considerable historical importance. Its significance stems from its association with Louis Agassiz, one of the principal figures in the introduction of the glacial theory in Scotland. The striated rock surface was unequivocally attributed by Agassiz to the effects of the passage of glacier ice. Agassiz Rock was also one of the first geological sites in Scotland to be conserved.

## Reference list

- Agassiz, L. (1841a) On glaciers and boulders in Switzerland. *Report of the British Association for 1840*, 113–4.
- Anon (1840) Tenth Meeting of the British Association for the Advancement of Science. Section C. – Geology and physical geography. *The Athenaeum*, No. **677**, 823–4.
- Brown, D.J. (1874) On some of the glacial phenomena of the neighbourhood of Edinburgh, as observed in the Pentlands, Blackford Hill, Bruntsfield Links, and Tynecastle Sandpit. *Transactions of the Edinburgh Geological Society*, **2**, 351–7.
- Buckland, W. (1841a) On the evidences of glaciers in Scotland and the north of England. *Proceedings of the Geological Society of London*, **3**, 332–7.
- Burke, M.J. (1968) Some stone-orientation results from the Forth valley. *Scottish Journal of Geology*, **5**, 286–92.
- Burke, M.J. (1969) The Forth valley: an ice-moulded lowland. *Transactions of the Institute of British Geographers*, **48**, 51–9.
- Campbell, R. (1951) Geology. In *Scientific Survey of South-eastern Scotland* Edinburgh, British Association, 184–99.
- Campbell, A.C. and Anderson, E.M. (1909) Notes on a transported mass of igneous rock at Comiston sand-pit, near Edinburgh. *Transactions of the Edinburgh Geological Society*, **9**, 219–24.
- Chambers, R. (1853) On glacial phenomena in Scotland and parts of England. *Edinburgh New Philosophical Journal*, **54**, 229–81.
- Cox, R. and Nicol, J. (1869) *Select Writings, Political, Scientific, Topographical, and Miscellaneous, of the Late Charles Maclaren, FRSE*. 2 vols. Edinburgh, Edmonston and Douglas.
- Davies, G.L. (1968a) *The Earth in Decay. A History of British Geomorphology 1578–1878*. London, MacDonald Technical and Scientific, 390pp.
- Davies, G.L. (1968b) The tour of the British Isles made by Louis Agassiz in 1840. *Annals of Science*, **24**, 131–46.
- Fleming, J. (1847) Notes on the superficial strata of the neighbourhood of Edinburgh, concluded. *Proceedings of the Royal Society of Edinburgh*, **2**, 111–3.
- Fleming, J. (1859) *The Lithology of Edinburgh*. (Edited, with a Memoir, by the Rev. John Duns) Edinburgh, William P. Kennedy. Memoir pp.i-civ.; Lithology pp.1–102.
- Geikie, A. (1863a) On the glacial drift of Scotland. *Transactions of the Geological Society of Glasgow*, **1**, 1–190.
- Geikie, A. (1901) *The Scenery of Scotland Viewed in Connection with its Physical Geology* 3rd edn. London, Macmillan and Co. 540pp.

- Goodchild, J.G. (1896) Glacial furrows. *The Glacialists' Magazine*, **4**, 1–7.
- Hall, Sir J. (1815) On the revolutions of the Earth's surface. *Transactions of the Royal Society of Edinburgh*, **7**, 139–211.
- Henderson, J. (1872) On Corstorphine Hill, near Edinburgh. *Transactions of the Edinburgh Geological Society*, **2**, 29–33.
- Henderson, J. (1874) On some sections of boulder clay, peat, and stratified beds, exposed in a quarry recently opened at Redhill, Slateford, near Edinburgh. *Transactions of the Edinburgh Geological Society*, **2**, 391–5.
- Imrie, (1812) A geological account of the southern district of Stirlingshire, commonly called the Campsie Hills, with a few remarks relative to the two prevailing theories as to Geology, and some examples given illustrative of these remarks. *Memoirs of the Wernerian Society*, **2**, 24–50.
- Kirby, R.P. (1969b) Till fabric analyses from the Lothians, central Scotland. *Geografiska Annaler*, **51A**, 48–60.
- Maclaren, C. (1828) Changes on the surface of the globe. *The Scotsman*, **12(918)**, 25 October, 683.
- Maclaren, C. (1840) Discovery of the former existence of glaciers in Scotland especially in the Highlands, by Professor Agassiz. *The Scotsman*, **24(2165)**, 7 October, 3.
- Maclaren, C. (1841) The glacial theory. No. 2. *The Scotsman*, **25(2190)**, 2 January, p.2.
- Maclaren, C. (1842a) The glacial theory of Prof Agassiz. *American Journal of Science and Arts*, **42**, 346–65.
- Maclaren, C. (1842b) Striated rocks of Corstorphine Hill. *The Scotsman*, **26(2344)**, 25 June, p.2.
- Maclaren, C. (1849) On grooved and striated rocks in the middle region of Scotland. *Edinburgh New Philosophical Journal*, **47**, 161–82.
- Maclaren, C. (1866) *A Sketch of the Geology of Fife and the Lothians Including Detailed Descriptions of Arthur's Seat and Pentland Hills*. Edinburgh, Adam and Charles Black, 2nd edition. 320pp.
- Miller, H. (1864) *Edinburgh and Its Neighbourhood, Geological and Historical, with the Geology of the Bass Rock*. Edinburgh, Adam and Charles Black, 313pp and 24pp.
- Miller, H. (1884) On boulder-glaciation. *Proceedings of the Royal Physical Society of Edinburgh* **8**, 156–189.
- Milne Home, D. (1840) On the Mid-Lothian and East-Lothian coal-fields. *Transactions of the Royal Society of Edinburgh*, **14**, 253–358.
- Milne Home, D. (1846) Notice of polished and striated rocks recently discovered on Arthur Seat, and in some other places near Edinburgh. *Edinburgh New Philosophical Journal*, **41**, 206–8.
- Milne Home, D. (1847a) On polished and striated rocks, lately discovered on Arthur Seat, and other places near Edinburgh. *Edinburgh New Philosophical Journal*, **42**, 154–72.
- Milne Home, D. (1871) *The Estuary of the Forth and Adjoining Districts Viewed Geologically*. Edinburgh, Edmonston and Douglas, 126 pp.
- Milne Home, D. (1874a) Notice of a striated boulder lately found in a sand pit at Tynecastle, near Edinburgh. *Transactions of the Edinburgh Geological Society*, **2**, 347–50.
- Milne Home, D. (1874b) Notice of a striated boulder found at Drylaw, near Linton, East Lothian. *Transactions of the Edinburgh Geological Society*, **2**, 350–1.
- Mitchell, G.H. and Mykura, W. (1962) The geology of the neighbourhood of Edinburgh. (Explanation of One-Inch Sheet 32). 3rd edn. Memoirs of the Geological Survey of Scotland. Edinburgh, HMSO, 159pp.
- Nicol, J. (1848) Observations on the recent formations in the vicinity of Edinburgh. *Quarterly Journal of the Geological Society of London*, **5**, 20–5.
- Panton, G.A. (1873) Note on a striated and water-worn cliff at Blackford Hill, near Edinburgh, and on a sandhill there. *Transactions of the Edinburgh Geological Society*, **2**, 238–42.
- Peach, A.M. (1909) Boulder distribution from Lennoxton, Scotland. *Geological Magazine*, **46**, 26–31.
- Peach, B.N., Clough, C.T., Hinxman, L.W., Wilson, J.S.G., Crampton, C.B., Maufe, H.B. and Bailey, E.B., (1910a) The Geology of the Neighbourhood of Edinburgh. (Explanation of Sheet 32, with part of 31). 2nd edn. Memoirs of the Geological Survey of Scotland. Edinburgh, HMSO, 445pp.
- Rhind, W. (1836) *Excursions Illustrative of the Geology and Natural History of the Environs of Edinburgh*. 2nd edn. Edinburgh, Maclachlan and Stewart, 72pp.

- 
- Richardson, R. (1877a) On phenomena of glaciation exhibited by the rocks of Corstorphine Hill, near Edinburgh. *Transactions of the Edinburgh Geological Society*, **3**, 31–40.
- Richardson, R. (1877b) Notice of glaciated rock surfaces (displaying corals) near Bathgate, recently quarried away. *Transactions of the Edinburgh Geological Society*, **3**, 108–9.
- Sissons, J.B. (1971) The geomorphology of central Edinburgh. *Scottish Geographical Magazine*, **87**, 185–96.
- Waterston, C.D. (1960) City of Edinburgh. In *Edinburgh Geology. An Excursion Guide* (eds G.H. Mitchell, E.K. Walton and D. Grant). Edinburgh, Oliver and Boyd, 10–27.
- Watson, J.A. (1934) General history. In *History of the Edinburgh Geological Society*, (hon. ed. R. Campbell). Edinburgh, Edinburgh Geological Society, 5–15.